

# Family planning and child health care: the Peruvian case, 1996-2000

Marianna Battaglia, Nina Pallarés | April 6, 2020



*In 1996-2000, for the first time in Peru, surgical contraception was promoted among indigenous women, and sometimes imposed upon them. As Marianna Battaglia and Nina Pallarés show, its overall positive impacts on contraceptive use and child mortality conceal substantial heterogeneity by ethnicity.*

## **The PNSRPF (1996-2000)**


Over the period 1996-2000, Peru carried out an unusual family planning program called *Programa de Salud Reproductiva y Planificación Familiar* (PNSRPF). At the time, infant and child mortality rates were exceptionally high in the country (45.8 and 23.5 per thousand live births in 1994, respectively) and there was a high prevalence of chronic malnutrition among children under five years (25.8% in 1996; UNICEF 2008). The Peruvian Ministry of Health promoted the PNSRPF with the stated purpose of addressing widespread poverty in the country through the reduction of fertility and of maternal and perinatal mortality. Its introduction was gradual, mainly through the celebration of health festivals that lasted about two days (Aramburú 2002). During these festivals, mobile medical teams visited the villages to provide information on family planning and contraception, and to offer health care services. Depending on the village health infrastructure, interventions occurred in the local health clinic or in the nearest hospital, or inside specially equipped ambulances (mobile sterilization units). All health services were provided free of charge, including voluntary surgical contraception, previously illegal.

However, several sources reported irregularities during the program implementation and documented episodes of forced sterilization among indigenous women from rural areas (Boesten 2007; Byker and Gutierrez 2012; Tamayo 1999). They were often pressured to have surgery through harassment of the household, and/or were offered money or food in exchange. The government administration refused to recognize such irregularities, never publicly stated

guidelines about target populations, and did not officially report the existence of sterilization quotas.

The peak years of the family planning campaign were 1996 and 1997. In 1998, the Committee of Latin America and the Caribbean for the Defense of Women's Rights (CLADEM) made a public statement accusing the family planning program of forced sterilization, and a first international investigation was launched. The Peruvian Ministry of Health replied to the accusations and changed the way the program goals were stated (Aramburú, 2002). Figure 1 reports the annual percentage of sterilizations among nonindigenous and indigenous women aged 15-49. DHS data show that indigenous women were less prone to choose sterilization both before and after the family planning program, in contrast with nonindigenous women who were already using permanent contraceptive methods even before it.

**Figure 1:** Percent of sterilizations per year for indigenous and nonindigenous women aged 15-49. DHS data.



## Consequences of the policy on women's use of contraceptive methods and newborn child health care

In a recent paper (Battaglia and Pallarés (2020)), we studied the effects of exposure to this unusual family planning program on child mortality and child health care and document differential child health care behaviors correlated with permanent or temporary interventions to reduce fertility, especially for those women among whom permanent intervention was aggressively promoted. To identify the areas of the country affected by the policy, we used the incidence and timing of surgical contraception among indigenous women in each province using data collected by the CLADEM (Tamayo 1999) and the Demographic and Health Surveys (DHS). We compared provinces where it is more likely that mobile medical units arrived earlier to celebrate health festivals (treated) with provinces where they arrived later (control),

and observed the impact of the policy in the short term, i.e. in the year when the policy reached the treated provinces.

As reported in Panel A of Table 1, due to the family planning program, women in treated areas were more likely to use any method of contraception (almost 10 percentage points), than women in control areas. The increase in sterilization was significantly higher (8.3 percentage points) for indigenous than for nonindigenous women.

Furthermore, children in treated provinces, born or conceived before the policy, were less likely to die within their first year of life compared to children in control provinces. Neonatal and infant mortality decreased by 5 to 6 percentage points (Panel B).

These results can be partially explained by a longer duration of breastfeeding in treated provinces (Panel C), where mothers were 18.5 percentage points more likely to breastfeed their children for longer than expected, given their age. On average, duration of breastfeeding increased by three weeks, from 10.2 to almost 11 months in this group. Breastmilk has immunological benefits and breastfeeding protects against water- and food-borne disease. Breastfeeding is also associated with lower rates of infant mortality from diarrheal disease and acute respiratory infection (Jayachandran and Kuziemko 2011).

**Table 1: Impacts of the PNSRPF**

	Treated (1)	Treated Indigenous (2)	Mean in control	Obs
<i>Panel A: Contraceptive methods</i>				
Any	0.099*** (0.025)	0.041 (0.046)	0.637	3434
Temporary	0.049*** (0.018)	-0.041 (0.033)	0.310	3434
Permanent	0.051*** (0.016)	0.083*** (0.027)	0.053	3434
<i>Panel B: Mortality and delivery</i>				
Neonatal <sup>(a)</sup>	-0.052*** (0.010)	-0.014 (0.017)	0.033	8688
Infant <sup>(b)</sup>	-0.061*** (0.013)	-0.003 (0.024)	0.058	7556
Home delivery	0.027 (0.030)	-0.052 (0.047)	0.588	2080
<i>Panel C: Breastfeeding and vaccines</i>				
Breastfeeding	0.185*** (0.060)	-0.131 (0.114)	0.791	1567
Vaccines at birth	-0.027 (0.062)	0.101 (0.133)	0.342	2477
Vaccines by 1 <sup>st</sup> year	0.021 (0.063)	0.021 (0.127)	0.246	1511

Robust standard errors clustered at the province level in parentheses: \* p < .10, \*\* p < .05, \*\*\* p < .01.

<sup>(a)</sup> In the first 28 days of life. <sup>(b)</sup> in the first year of life.

Column (1) reports the effect of treated areas versus control areas; column (2) of indigenous versus nonindigenous in treated areas only. All the dependent variables are dummies and the corresponding coefficients must be interpreted as relative variations from the baseline value of the control group (mean in control).

Medical research also shows that childhood vaccination, in particular against measles and tetanus, is associated with substantial reductions in child mortality (World Health Organization 2017). However, children's likelihood of receiving the appropriate vaccinations at birth and by the first year of life was not significantly affected by the policy.

By comparing women in treated provinces who used a contraceptive method with those who did not, we observe that nonindigenous children whose mothers used contraception benefited from the policy whatever the method used (Table 2). The children of nonindigenous women who adopted temporary methods were 17.6 percentage points less likely to die in their the first year of life and 7.1 percentage points less likely to be delivered at home, compared to children whose mothers did not use any contraceptive method. The children of sterilized mothers were 23.1 percentage points more likely to be breastfed for longer.

**Table 2: Treated provinces: contraceptive methods on mortality, breastfeeding and vaccination**

	Indigenous		Mean in control	Obs	Nonindigenous		Mean in control	Obs
	Temporary (1)	Permanent (2)			Temporary (3)	Permanent (4)		
<i>Panel A: Mortality and delivery</i>								
Neonatal	0.046 (0.057)	-0.019* (0.011)	0.018	618	0.004 (0.003)	0.008 (0.006)	0.007	445
Infant	-0.030 (0.046)	-0.082*** (0.017)	0.074	906	-0.070* (0.035)	-0.026 (0.025)	0.048	722
Home delivery	-0.003 (0.073)	-0.303** (0.132)	0.884	233	-0.076** (0.069)	-0.239 (0.200)	0.503	159
<i>Panel B: Breastfeeding and vaccines</i>								
Breastfeeding	0.168*** (0.043)	0.001 (0.221)	0.752	399	0.042 (0.046)	0.231* (0.123)	0.173	300
Vaccines at birth	0.261** (0.098)	-0.055 (0.170)	0.330	312	0.168 (0.1116)	-0.135 (0.234)	0.387	191
Vaccines by 1st year	0.143 (0.092)	-0.010 (0.125)	0.260	288	0.135 (0.097)	0.060 (0.171)	0.310	184

Robust standard errors clustered at the province level in parentheses: \*  $p < .10$ , \*\*  $p < .05$ , \*\*\*  $p < .01$ .  
How to read the table: See note to Table (1)

Conversely, among indigenous mothers, the effects on their children's health care were positive if they used temporary methods but much less so if they were sterilized. Children whose mothers used temporary contraception were both more likely to be breastfed for longer (roughly one month more) and to receive the required vaccinations at birth (26.1 percentage points), compared to children whose mothers did not use any contraception. Even though mortality was lower among children born to mothers who were sterilized, this seems to be mainly related to safer conditions of childbirth rather than better child care after delivery.

## What did we learn?

DHS data do not tell us who freely decided to adopt a permanent contraceptive method and who was forced to do so. Therefore, we cannot derive strong conclusions and claim any causal interpretation of the impact of contraception on child health. Nevertheless, these results are informative of differences by ethnicity in reactions to the policy and of the heterogeneous impacts on child health care when a policy intervention is aggressively promoted. As Byker and Gutierrez (2012) noted, when birth control is imposed, the benefits of making choices about fertility are lost and having fewer children may not translate into substantial improvements in the health care they receive.

## References

Aramburú, Carlos E. 2002. "Politics and Reproductive Health: A Dangerous Connection." in *Interregional Seminar on Reproductive Health, Unmet Needs and Poverty*. Bangkok, Thailand.

Paper presented at the CICRED conference, 25-29 Nov.

Battaglia, Marianna and Nina Pallarés. 2020. "Family Planning and Child Health Care: Effect of the Peruvian Programa de Salud Reproductiva y Planificación Familiar, 1996- 2000". *Population and Development Review*, forthcoming.

Boesten, Jelke. 2007. "Free Choice or Poverty Alleviation? Population Politics in Peru under Alberto Fujimori." *European Review of Latin American and Caribbean Studies* 82, 3-20.

Byker, Tanya and Italo A. Gutierrez. 2012. "Fertility and Family Well-being Effects of an Aggressive Family Planning Policy in Peru in the 1990s." *PSC Research Report* No. 12-765.

Jayachandran, Seema and Ilyana Kuziemko. 2011. "Why Do Mothers Breastfeed Girls Less than Boys: Evidence and Implications for Child Health in India." *Quarterly Journal of Economics* 12 (3), 1485-1538.

Tamayo, Julia. 1999. "Nada Personal: Reporte de Derechos Humanos Sobre la Aplicación de la Anticoncepción Quirúrgica en el Perú, 1996-1998." Technical report, Comité de América Latina y el Caribe para la Defensa de los Derechos de la Mujer (CLADEM), Lima.

UNICEF. 2008. Situation of Children in Peru. Executive Summary. Technical report, United Nations Children's Fund (UNICEF) and Instituto Nacional de Estadística e Informática (INEI).

World Health Organization. 2017. 2017 Assessment Report of the Global Vaccine Action Plan. Technical report, Strategic Advisory Group of Experts on Immunization on the Progress towards GVAP Targets. World Health Organization 2017.